

Hadron Form Factors and DSEs

Craig D. Roberts


cdroberts@anl.gov

Physics Division
Argonne National Laboratory

<http://www.phy.anl.gov/theory/staff/cdr.html>


Universal Truths

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Universal Truths


[First](#)[Contents](#)[Back](#)[Conclusion](#)



Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe.





Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe. Higgs mechanism is irrelevant to light-quarks.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe. Higgs mechanism is irrelevant to light-quarks.
- Challenge: understand relationship between parton properties on the light-front and rest frame structure of hadrons.





Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe. Higgs mechanism is irrelevant to light-quarks.
- Challenge: understand relationship between parton properties on the light-front and rest frame structure of hadrons. Problem because, e.g., DCSB - an established keystone of low-energy QCD and the origin of constituent-quark masses - has not been realised in the light-front formulation.



QCD's Challenges




[First](#)

[Contents](#)


[Back](#)

[Conclusion](#)




- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon





- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon
- Dynamical Chiral Symmetry Breaking
 - Very unnatural pattern of bound state masses
 - e.g., Lagrangian (pQCD) quark mass is small but ... no degeneracy between $J^{P=+}$ and $J^{P=-}$





- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon
- Dynamical Chiral Symmetry Breaking
 - Very unnatural pattern of bound state masses
 - e.g., Lagrangian (pQCD) quark mass is small but ... no degeneracy between $J^{P=+}$ and $J^{P=-}$
- Neither of these phenomena is apparent in QCD's Lagrangian **yet** they are the dominant determining characteristics of real-world QCD.



Understand Emergent Phenomena

- Quark and Gluon Confinement
 - No matter how hard one strikes the proton, one cannot liberate an individual quark or gluon
- Dynamical Chiral Symmetry Breaking
 - Very unnatural pattern of bound state masses
 - e.g., Lagrangian (pQCD) quark mass is small but ... no degeneracy between $J^{P=+}$ and $J^{P=-}$
- Neither of these phenomena is apparent in QCD's Lagrangian **yet** they are the dominant determining characteristics of real-world QCD.
- QCD – Complex behaviour
arises from apparently simple rules



Why?

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.



Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.



Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.
- Form factors have long been recognized as a basic tool for elucidating bound state properties. They can be studied from very low momentum transfer, the region of non-perturbative QCD, up to a region where perturbative QCD predictions can be tested.



Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.
- Form factors have long been recognized as a basic tool for elucidating bound state properties. They can be studied from very low momentum transfer, the region of non-perturbative QCD, up to a region where perturbative QCD predictions can be tested.
- Experimental and theoretical studies of nucleon electromagnetic form factors have made rapid and significant progress during the last several years, including new data in the time like region, and material gains have been made in studying the pion form factor.



Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.
- Form factors have long been recognized as a basic tool for elucidating bound state properties. They can be studied from very low momentum transfer, the region of non-perturbative QCD, up to a region where perturbative QCD predictions can be tested.
- Experimental and theoretical studies of nucleon electromagnetic form factors have made rapid and significant progress during the last several years, including new data in the time like region, and material gains have been made in studying the pion form factor.
- Despite this, many urgent questions remain unanswered.



Some Questions

- What is the role of pion cloud in nucleon electromagnetic structure?
- Can we understand the pion cloud in a more quantitative and, perhaps, model-independent way?



Some Questions

- Where is the transition from non-pQCD to pQCD in the pion and nucleon electromagnetic form factors?



Some Questions

- Do we understand the high Q^2 behavior of the proton form factor ratio in the space-like region?
- Can we make model-independent statements about the role of relativity or orbital angular momentum in the nucleon?



Some Questions

- Can we understand the rich structure of the time-like proton form factors in terms of resonances?
- What do we expect for the proton form factor ratio in the time-like region?
- What is the relation between proton and neutron form factor in the time-like region?
- How do we understand the ratio between time-like and space-like form factors?



Some Questions

- What is the role of two-photon exchange contributions in understanding the discrepancy between the polarization and Rosenbluth measurements of the proton form factor ratio?
- What is the impact of these contributions on other form factor measurements?



Some Questions

- How accurately can the pion form factor be extracted from the $ep \rightarrow e'n\pi^+$ reaction?





- Current status is described in
 - J. Arrington, C. D. Roberts and J. M. Zanotti
“Nucleon electromagnetic form factors,”
J. Phys. G **34**, S23 (2007); [arXiv:nucl-th/0611050].
 - C. F. Perdrisat, V. Punjabi and M. Vanderhaeghen,
“Nucleon electromagnetic form factors,”
Prog. Part. Nucl. Phys. **59**, 694 (2007);
[arXiv:hep-ph/0612014].



- Current status is described in
 - J. Arrington, C. D. Roberts and J. M. Zanotti
“Nucleon electromagnetic form factors,”
J. Phys. G **34**, S23 (2007); [arXiv:nucl-th/0611050].
 - C. F. Perdrisat, V. Punjabi and M. Vanderhaeghen,
“Nucleon electromagnetic form factors,”
Prog. Part. Nucl. Phys. **59**, 694 (2007);
[arXiv:hep-ph/0612014].

- Most recently:
“ECT* Workshop on Hadron Electromagnetic Form Factors”
Organisers: Alexandrou, Arrington, Friedrich, Maas, Roberts
Presentations, etc., available on-line
<http://ect08.phy.anl.gov/>



Dichotomy of Pion

– Goldstone Mode and Bound state





Dichotomy of Pion

– Goldstone Mode and Bound state

- How does one make an **almost massless** particle
..... from two **massive** constituent-quarks?





Dichotomy of Pion

– Goldstone Mode and Bound state

- How does one make an **almost massless** particle
..... from two **massive** constituent-quarks?
- **Not Allowed** to do it by **fine-tuning** a potential

Must exhibit $m_\pi^2 \propto m_q$

Current Algebra ... 1968





Dichotomy of Pion

– Goldstone Mode and Bound state

- How does one make an **almost massless** particle from two **massive** constituent-quarks?
- **Not Allowed** to do it by **fine-tuning** a potential

Must exhibit $m_\pi^2 \propto m_q$

Current Algebra ... 1968

The **correct understanding** of pion observables; e.g. **mass**, **decay constant** and **form factors**, **requires** an approach to contain a

- **well-defined** and **valid chiral limit**;
- and an **accurate realisation** of dynamical chiral symmetry breaking.





Dichotomy of Pion

– Goldstone Mode and Bound state

- How does one make an **almost massless** particle from two **massive** constituent-quarks?
- **Not Allowed** to do it by **fine-tuning** a potential

Must exhibit $m_\pi^2 \propto m_q$

Current Algebra ... 1968

The **correct understanding** of pion observables; e.g. **mass**, **decay constant** and **form factors**, **requires** an approach to contain a

- **well-defined** and **valid chiral limit**;
- and an **accurate realisation** of dynamical chiral symmetry breaking.

Highly Nontrivial



Pion Form Factors

[First](#)[Contents](#)[Back](#)[Conclusion](#)

- There is a sense in which it is easy to fabricate a model that can *reproduce* the elastic electromagnetic pion form factor



- There is a sense in which it is easy to fabricate a model that can *reproduce* the elastic electromagnetic pion form factor
- However, a **veracious description** of the pion will **simultaneously predict** the elastic electromagnetic form factor, $F_\pi(Q^2)$ AND the $\gamma^* \pi \rightarrow \gamma$ transition form factor



Infidelity without simultaneity

- There is a sense in which it is easy to fabricate a model that can *reproduce* the elastic electromagnetic pion form factor
- However, a **veracious description** of the pion will **simultaneously predict** the elastic electromagnetic form factor, $F_\pi(Q^2)$ AND the $\gamma^* \pi \rightarrow \gamma$ transition form factor
- The latter is connected with the **Abelian anomaly** – therefore **fundamentally connected** with **chiral symmetry and its dynamical breaking** – no mere model can successfully describe this without fine tuning



Infidelity without simultaneity

- There is a sense in which it is easy to fabricate a model that can *reproduce* the elastic electromagnetic pion form factor
- However, a **veracious description** of the pion will **simultaneously predict** the elastic electromagnetic form factor, $F_\pi(Q^2)$ AND the $\gamma^* \pi \rightarrow \gamma$ transition form factor
- The latter is connected with the **Abelian anomaly** – therefore **fundamentally connected** with **chiral symmetry and its dynamical breaking** – no mere model can successfully describe this without fine tuning
- Must similarly require prediction of $\gamma^* \pi \rightarrow \pi\pi$ and all other anomalous processes



What's the Problem?

[First](#)[Contents](#)[Back](#)[Conclusion](#)

What's the Problem?

- Minimal requirements
 - detailed understanding of connection between **Current-quark** and **Constituent-quark** masses;
 - and systematic, symmetry preserving means of realising this connection in bound-states.



What's the Problem?

- Minimal requirements
 - detailed understanding of connection between **Current-quark** and **Constituent-quark** masses;
 - and systematic, symmetry preserving means of realising this connection in bound-states.
- Means ... must calculate hadron *wave functions*
 - Can't be done using perturbation theory



What's the Problem?

- Minimal requirements
 - detailed understanding of connection between **Current-quark** and **Constituent-quark** masses;
 - and systematic, symmetry preserving means of realising this connection in bound-states.
- Means ... must calculate hadron *wave functions*
 - Can't be done using perturbation theory
- Why problematic? Isn't same true in quantum mechanics?



What's the Problem?

- Minimal requirements
 - detailed understanding of connection between **Current-quark** and **Constituent-quark** masses;
 - and systematic, symmetry preserving means of realising this connection in bound-states.
- Means ... must calculate hadron *wave functions*
 - Can't be done using perturbation theory
- Why problematic? Isn't same true in quantum mechanics?
- Differences!



What's the Problem?

Relativistic QFT!

- Minimal requirements
 - detailed understanding of connection between **Current-quark** and **Constituent-quark** masses;
 - and systematic, symmetry preserving means of realising this connection in bound-states.
- Differences!
 - Here relativistic effects are crucial – *virtual particles*, quintessence of **Relativistic Quantum Field Theory** – must be included



What's the Problem?

Relativistic QFT!


- Minimal requirements
 - detailed understanding of connection between **Current-quark** and **Constituent-quark** masses;
 - and systematic, symmetry preserving means of realising this connection in bound-states.
- Differences!
 - Here relativistic effects are crucial – *virtual particles*, quintessence of **Relativistic Quantum Field Theory** – must be included
 - Interaction between quarks – the **Interquark “Potential”** – *unknown* throughout **> 98%** of a hadron's volume




Intranucleon Interaction

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Intranucleon Interaction



Intranucleon Interaction




98% of the volume



What is the Intranucleon Interaction?

The question must be rigorously defined, and the answer mapped out using experiment and theory.



98% of the volume



Dyson-Schwinger Equations

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Dyson-Schwinger Equations


Dressed-Quark Propagator



Dyson-Schwinger Equations

Dressed-Quark Propagator

$$S(p) = \frac{Z(p^2)}{i\gamma \cdot p + M(p^2)}$$




Gap Equation



Dyson-Schwinger Equations


Dressed-Quark Propagator

$$S(p) = \frac{Z(p^2)}{i\gamma \cdot p + M(p^2)}$$



Gap Equation


- Gap Equation's Kernel Enhanced on IR domain
- ⇒ IR Enhancement of $M(p^2)$



Dyson-Schwinger Equations

Dressed-Quark Propagator

$$S(p) = \frac{Z(p^2)}{i\gamma \cdot p + M(p^2)}$$




Gap Equation

- Gap Equation's Kernel Enhanced on IR domain
- ⇒ IR Enhancement of $M(p^2)$

Euclidean Constituent-Quark

Mass: $M_f^E: p^2 = M(p^2)^2$


flavour	u/d	s	c	b
$\frac{M^E}{m_\zeta}$	$\sim 10^2$	~ 10	~ 1.5	~ 1.1



Dyson-Schwinger Equations

Dressed-Quark Propagator

$$S(p) = \frac{Z(p^2)}{i\gamma \cdot p + M(p^2)}$$




Gap Equation

- Gap Equation's Kernel Enhanced on IR domain
- ⇒ IR Enhancement of $M(p^2)$

Euclidean Constituent-Quark

Mass: $M_f^E: p^2 = M(p^2)^2$

flavour	u/d	s	c	b
$\frac{M^E}{m_\zeta}$	$\sim 10^2$	~ 10	~ 1.5	~ 1.1




Predictions confirmed in numerical simulations of lattice-QCD





Frontiers of Nuclear Science: A Long Range Plan (2007)

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Frontiers of Nuclear Science: Theoretical Advances




Frontiers of Nuclear Science: Theoretical Advances



Frontiers of Nuclear Science: Theoretical Advances

Mass from nothing.


In QCD a quark's effective mass depends on its momentum. The function describing this can be calculated and is depicted here. Numerical simulations of lattice QCD (data, at two different bare masses) have confirmed model predictions (solid curves) that the vast bulk of the constituent mass of a light quark comes from a cloud of gluons that are dragged along by the quark as it propagates. In this way, a quark that appears to be absolutely massless at high energies ($m = 0$, red curve) acquires a large constituent mass at low energies.



Frontiers of Nuclear Science: Theoretical Advances

Mass from nothing.

In QCD a quark's effective mass depends on its momentum. The function describing this can be calculated and is depicted here. Numerical simulations of lattice QCD (data, at two different bare masses) have confirmed **model predictions (solid curves)** that the vast bulk of the constituent mass of a light quark comes from a cloud of gluons that are dragged along by the quark as it **propagates**. In this way, a quark that appears to be absolutely massless at high energies ($m = 0$, red curve) acquires a large constituent mass at low energies.





- Established understanding of two- and three-point functions





- Established understanding of two- and three-point functions
- What about bound states?



Hadrons




- Without bound states,
Comparison with experiment is
impossible



- Without bound states,
Comparison with experiment is
impossible
- They appear as pole contributions
to $n \geq 3$ -point colour-singlet
Schwinger functions




- Without bound states,
Comparison with experiment is
impossible
- Bethe-Salpeter Equation



QFT Generalisation of Lippmann-Schwinger Equation.


- Without bound states,
Comparison with experiment is
impossible
- Bethe-Salpeter Equation



QFT Generalisation of Lippmann-Schwinger Equation.

- What is the kernel, K ?

- Without bound states,
Comparison with experiment is
impossible
- Bethe-Salpeter Equation



QFT Generalisation of Lippmann-Schwinger Equation.

- What is the kernel, K ?
- or

What is the light-quark Long-Range Potential?



What is the light-quark Long-Range Potential?



Potential between static (infinitely heavy) quarks measured in simulations of lattice-QCD **is not related** in any simple way to the light-quark interaction.



Bethe-Salpeter Kernel

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-)$$

$$-M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

QFT Statement of Chiral Symmetry



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-) - M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE



Bethe-Salpeter Kernel

● Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-) - M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE

Kernels very different
but must be *intimately* related



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-) - M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE

Kernels very different
but must be *intimately* related

- Relation **must** be preserved by truncation



Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-) - M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE

Kernels very different
but must be *intimately* related

- Relation **must** be preserved by truncation
- **Nontrivial** constraint





Bethe-Salpeter Kernel

- Axial-vector Ward-Takahashi identity

$$P_\mu \Gamma_{5\mu}^l(k; P) = \mathcal{S}^{-1}(k_+) \frac{1}{2} \lambda_f^l i \gamma_5 + \frac{1}{2} \lambda_f^l i \gamma_5 \mathcal{S}^{-1}(k_-) - M_\zeta i \Gamma_5^l(k; P) - i \Gamma_5^l(k; P) M_\zeta$$

Satisfies BSE

Satisfies DSE

Kernels very different

but must be *intimately* related

- Relation **must** be preserved by truncation
- **Failure** \Rightarrow Explicit Violation of QCD's Chiral Symmetry





Persistent Challenge

- Infinitely Many Coupled Equations
- There is at least one **systematic nonperturbative, symmetry-preserving** truncation scheme

H.J. Munczek Phys. Rev. D **52** (1995) 4736

Dynamical chiral symmetry breaking, Goldstone's theorem and the consistency of the Schwinger-Dyson and Bethe-Salpeter Equations

A. Bender, C. D. Roberts and L. von Smekal, Phys. Lett. B **380** (1996) 7

Goldstone Theorem and Diquark Confinement Beyond Rainbow Ladder Approximation





Persistent Challenge

- Infinitely Many Coupled Equations
- There is at least one **systematic nonperturbative, symmetry-preserving** truncation scheme
- Has Enabled Proof of **EXACT** Results in QCD





Persistent Challenge

- Infinitely Many Coupled Equations
- There is at least one **systematic nonperturbative, symmetry-preserving** truncation scheme
- Has Enabled Proof of **EXACT** Results in QCD
- And Formulation of Practical Phenomenological Tool to
 - Illustrate Exact Results





Persistent Challenge

- Infinitely Many Coupled Equations
- There is at least one **systematic nonperturbative, symmetry-preserving** truncation scheme
- Has Enabled Proof of **EXACT** Results in QCD
- And Formulation of Practical Phenomenological Tool to
 - Illustrate Exact Results
 - Make Predictions with Readily Quantifiable Errors



Radial Excitations & Chiral Symmetry

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \, m_H^2 = - \, \rho_{\zeta}^H \, \mathcal{M}_H$$



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \, m_H^2 = - \, \rho_\zeta^H \, \mathcal{M}_H$$

- Mass² of pseudoscalar hadron



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \, m_H^2 = - \, \rho_\zeta^H \, \mathcal{M}_H$$

$$\mathcal{M}_H := \text{tr}_{\text{flavour}} \left[M_{(\mu)} \left\{ T^H, (T^H)^t \right\} \right] = m_{q_1} + m_{q_2}$$

- Sum of constituents' current-quark masses
- e.g., $T^{K^+} = \frac{1}{2} (\lambda^4 + i\lambda^5)$




Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

$$f_H p_\mu = Z_2 \int_q^\Lambda \frac{1}{2} \text{tr} \left\{ (T^H)^t \gamma_5 \gamma_\mu \mathcal{S}(q_+) \Gamma_H(q; P) \mathcal{S}(q_-) \right\}$$

- Pseudovector projection of BS wave function at $x = 0$
- Pseudoscalar meson's leptonic decay constant




Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H \quad m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

$$i\rho_\zeta^H = Z_4 \int_q^\Lambda \frac{1}{2} \text{tr} \left\{ (T^H)^t \gamma_5 \mathcal{S}(q_+) \Gamma_H(q; P) \mathcal{S}(q_-) \right\}$$

- Pseudoscalar projection of BS wave function at $x = 0$



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

● Light-quarks; i.e., $m_q \sim 0$

● $f_H \rightarrow f_H^0$ & $\rho_\zeta^H \rightarrow \frac{-\langle \bar{q}q \rangle_\zeta^0}{f_H^0}$, Independent of m_q

Hence $m_H^2 = \frac{-\langle \bar{q}q \rangle_\zeta^0}{(f_H^0)^2} m_q \dots$ GMOR relation, a corollary



Radial Excitations & Chiral Symmetry

(Maris, Roberts, Tandy
nu-th/9707003)

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

- Light-quarks; i.e., $m_q \sim 0$

- $f_H \rightarrow f_H^0$ & $\rho_\zeta^H \rightarrow \frac{-\langle \bar{q}q \rangle_\zeta^0}{f_H^0}$, Independent of m_q

Hence $m_H^2 = \frac{-\langle \bar{q}q \rangle_\zeta^0}{(f_H^0)^2} m_q \dots$ GMOR relation, a corollary

- Heavy-quark + light-quark

$\Rightarrow f_H \propto \frac{1}{\sqrt{m_H}}$ and $\rho_\zeta^H \propto \sqrt{m_H}$

Hence, $m_H \propto m_q$

\dots QCD Proof of Potential Model result



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson, not the ground state, so $m_{\pi_{n \neq 0}}^2 > m_{\pi_{n=0}}^2 = 0$, in chiral limit



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson, not the ground state, so $m_{\pi_{n \neq 0}}^2 > m_{\pi_{n=0}}^2 = 0$, in chiral limit
- $\Rightarrow f_H = 0$
ALL pseudoscalar mesons except $\pi(140)$ in chiral limit



Radial Excitations & Chiral Symmetry

Höll, Krassnigg, Roberts
nu-th/0406030

$$f_H m_H^2 = - \rho_\zeta^H \mathcal{M}_H$$

- Valid for ALL Pseudoscalar mesons
- $\rho_H \Rightarrow$ finite, nonzero value in chiral limit, $\mathcal{M}_H \rightarrow 0$
- “radial” excitation of π -meson, not the ground state, so $m_{\pi_{n \neq 0}}^2 > m_{\pi_{n=0}}^2 = 0$, in chiral limit
- $\Rightarrow f_H = 0$
ALL pseudoscalar mesons except $\pi(140)$ in chiral limit
- Dynamical Chiral Symmetry Breaking
 - Goldstone’s Theorem –impacts upon every pseudoscalar meson



Pion Form Factor

Procedure Now Straightforward



[First](#)


[Contents](#)

[Back](#)

[Conclusion](#)


Pion Form Factor

- Solve Gap Equation
 - ⇒ Dressed-Quark Propagator, $S(p)$




Pion Form Factor

- Use that to Complete Bethe Salpeter Kernel, K
- Solve Homogeneous Bethe-Salpeter Equation for Pion Bethe-Salpeter Amplitude, Γ_π



Pion Form Factor

- Use that to Complete Bethe Salpeter Kernel, K
- Solve Homogeneous Bethe-Salpeter Equation for Pion Bethe-Salpeter Amplitude, Γ_π




- Solve Inhomogeneous Bethe-Salpeter Equation for Dressed-Quark-Gluon Vertex, Γ_μ




Pion Form Factor

- Now have all elements for Impulse Approximation to Electromagnetic Pion Form factor



Pion Form Factor

- Now have all elements for Impulse Approximation to Electromagnetic Pion Form factor




- Evaluate this final, three-dimensional integral



Calculated Pion Form Factor

Calculation first published in 1999; No Parameters Varied

Numerical method improved in 2005




Calculated Pion Form Factor

Calculation first published in 1999; No Parameters Varied

Numerical method improved in 2005

Data published
in 2001.
Subsequently
revised





Timelike Pion Form Factor

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Timelike Pion Form Factor


Ab initio calculation into timelike region
Deeper than ground-state ρ -meson pole





Timelike Pion Form Factor

Ab initio calculation into timelike region
Deeper than ground-state ρ -meson pole



Office of Science
U.S. DEPARTMENT OF ENERGY

Office of Nuclear Physics
Exploring Nuclear Matter - Quarks in Stars



Argonne
NATIONAL
LABORATORY

First

Contents

Back

Cor




Timelike Pion Form Factor

Ab initio calculation into timelike region

Deeper than ground-state ρ -meson pole

ρ -meson not put in “by hand” – generated dynamically as a bound-state of dressed-quark and dressed-antiquark



Dimensionless product: $r_\pi f_\pi$





Dimensionless product: $r_\pi f_\pi$

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction





Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation





Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Great strides towards placing nucleon studies on same footing as mesons



Dimensionless product: $r_\pi f_\pi$


- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$



Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$

DSE prediction




Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$

DSE prediction

Lattice results


– James Zanotti [UK QCD]



Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$


- DSE prediction
- Lattice results
 - James Zanotti [UK QCD]
- Fascinating result:
DSE and Lattice
 - Experimental value obtains independent of current-quark mass.



Dimensionless product: $r_\pi f_\pi$


- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$

- DSE prediction
- Fascinating result:
DSE and Lattice
– Experimental value
obtains independent of
current-quark mass.
We have understood this
Implications far-reaching.



Two-photon Couplings of Pseudoscalar Mesons

Höll, Krassnigg, Maris, *et al.*,
 “Electromagnetic properties of ground and
 excited state pseudoscalar mesons,”
 nu-th/0503043



$$T_{\mu\nu}^{\pi_n^0}(k_1, k_2) = \frac{\alpha}{\pi} i \varepsilon_{\mu\nu\rho\sigma} k_{1\rho} k_{2\sigma} G^{\pi_n^0}(k_1, k_2)$$

Define: $\mathcal{T}_{\pi_n^0}(P^2, Q^2) = G^{\pi_n^0}(k_1, k_2) \Big|_{k_1^2 = Q^2 = k_2^2}$

This is a transition form factor.

Physical Processes described by couplings:


$$g_{\pi_0^0 \gamma \gamma} := \mathcal{T}_{\pi_0^0}(-m_{\pi_0^0}^2, 0)$$

$$\text{Width: } \Gamma_{\pi_n^0 \gamma \gamma} = \alpha_{\text{em}}^2 \frac{m_{\pi_n}^3}{16\pi^3} g_{\pi_n \gamma \gamma}^2$$



Two-photon Couplings: Goldstone Mode

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043



$$T_{\mu\nu}^{\pi_0^0}(k_1, k_2) = \frac{\alpha}{\pi} i \varepsilon_{\mu\nu\rho\sigma} k_{1\rho} k_{2\sigma} G^{\pi_0^0}(k_1, k_2)$$

Chiral limit, model-independent and algebraic result

$$g_{\pi_0^0\gamma\gamma} := \mathcal{T}_{\pi_0^0}(-m_{\pi_0^0}^2 = 0, 0) = \frac{1}{2} \frac{1}{f_{\pi_0}}$$

So long as truncation veraciously preserves chiral symmetry
and the pattern of its dynamical breakdown


The most widely known consequence of the **Abelian anomaly**



Argonne
NATIONAL
LABORATORY

Two-photon Couplings: Transition Form Factor

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043



$$T_{\mu\nu}^{\pi_n^0}(k_1, k_2) = \frac{\alpha}{\pi} i \varepsilon_{\mu\nu\rho\sigma} k_{1\rho} k_{2\sigma} G^{\pi_n^0}(k_1, k_2)$$


So long as truncation preserves **chiral symmetry** and the pattern of its dynamical breakdown, and the one-loop renormalisation group properties of QCD: model-independent result – $\forall n$:

$$\mathcal{T}_{\pi_n^0}(P^2, Q^2) = G^{\pi_n^0}(k_1, k_2) \Big|_{k_1^2 = Q^2 = k_2^2} \stackrel{Q^2 \gg \Lambda_{\text{QCD}}^2}{=} \frac{4\pi^2}{3} \frac{f_{\pi_n}}{Q^2}$$



Two-photon Couplings: Transition Form Factor

Maris and Tandy, " Electromagnetic
transition form-factors of light mesons,"
nucl-th/0201017




Two-photon Couplings: Transition Form Factor

Maris and Tandy, " Electromagnetic
transition form-factors of light mesons,"
nucl-th/0201017

DSE result:

- normalisation
calculated
- ρ -meson generated
dynamically
- pQCD result accurate
to $\sim 20\%$ or better for
 $Q^2 \geq 3 \text{ GeV}^2$



Transition Form Factor: Chiral limit

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043

● Chiral limit with DCSB: $f_{\pi_0} \neq 0$



Transition Form Factor:

Chiral limit

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043

● Chiral limit with DCSB: $f_{\pi_0} \neq 0$

● **BUT**, $f_{\pi_n} \equiv 0, \forall n!$



Transition Form Factor: Chiral limit

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043

- Chiral limit with DCSB: $f_{\pi_0} \neq 0$
- **BUT**, $f_{\pi_n} \equiv 0, \forall n!$
- Model-independent result, in chiral limit: $\forall n \geq 1$

$$\lim_{\hat{m} \rightarrow 0} \mathcal{T}_{\pi_n^0}(-m_{\pi_n}^2, Q^2)$$

$$\stackrel{Q^2 \gg \Lambda_{\text{QCD}}^2}{=} \frac{4\pi^2}{3} F_n^{(2)}(-m_{\pi_n}^2) \frac{\ln^\gamma Q^2 / \omega_{\pi_n}^2}{Q^4} \Big|_{\hat{m}=0}$$

where:

- γ is an anomalous dimension
- ω_{π_n} is a width mass-scale

both determined, in part, by properties of the meson's
Bethe-Salpeter wave function.



Höll, Krassnigg, Maris, *et al.*,
 “Electromagnetic properties of ground and
 excited state pseudoscalar mesons,”
 nu-th/0503043

- Chiral limit with DCSB: $f_{\pi_0} \neq 0$
- **BUT**, $f_{\pi_n} \equiv 0, \forall n!$
- Model-independent result, in chiral limit: $\forall n \geq 1$

$$\lim_{\hat{m} \rightarrow 0} \mathcal{T}_{\pi_n^0}(-m_{\pi_n}^2, Q^2)$$

$$\stackrel{Q^2 \gg \Lambda_{\text{QCD}}^2}{=} \frac{4\pi^2}{3} F_n^{(2)}(-m_{\pi_n}^2) \frac{\ln^\gamma Q^2 / \omega_{\pi_n}^2}{Q^4} \Big|_{\hat{m}=0}$$

where:

- γ is an anomalous dimension
- ω_{π_n} is a width mass-scale

both determined, in part, by properties of the meson's
 Bethe-Salpeter wave function.


- Importantly, $F_n^{(2)}(-m_{\pi_n}^2) \not\propto f_{\pi_n}$. Instead, it is determined by
 DCSB mass-scales for π_n that do not vanish in the chiral limit.




Transition Form Factor (*Chiral*):

RGI Rainbow-Ladder

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043




 $m_u(1 \text{ GeV})$
 $= m_d(1 \text{ GeV}) = 0$



Transition Form Factor (*Chiral*):

RGI Rainbow-Ladder

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043




- $m_u(1 \text{ GeV}) = m_d(1 \text{ GeV}) = 0$
- Again, Predicted *UV*-behaviour is abundantly clear
 - precise for $Q^2 > 120 \text{ GeV}^2$



Transition Form Factor (*Chiral*):

RGI Rainbow-Ladder

Höll, Krassnigg, Maris, *et al.*,
 “Electromagnetic properties of ground and
 excited state pseudoscalar mesons,”
 nu-th/0503043



- $m_u(1 \text{ GeV})$
 $= m_d(1 \text{ GeV}) = 0$

- Again, Predicted *UV*-behaviour
 is abundantly clear

- precise for $Q^2 > 120 \text{ GeV}^2$

- $$F_1^{(2)}(-m_{\pi_1}^2) \ln^\gamma Q^2 / \omega_{\pi_1}^2 \Big|_{\hat{m}=0} \approx (0.22 \text{ GeV})^3 \simeq -\langle \bar{q}q \rangle^0 \quad (3)$$




Nucleon Challenge

[First](#)[Contents](#)[Back](#)[Conclusion](#)


Nucleon Challenge

- Another Direction . . . Also want/need information about three-quark systems




Nucleon Challenge


- Another Direction . . . Also want/need information about three-quark systems
- With this problem . . . current expertise at approximately same point as studies of mesons in 1995.



Nucleon Challenge

- Another Direction . . . Also want/need information about three-quark systems
- With this problem . . . current expertise at approximately same point as studies of mesons in 1995.
- Namely . . . Model-building and Phenomenology, constrained by the DSE results outlined already.





Nucleon EM Form Factors: A Précis

Höll, et al.: nu-th/0412046 & nu-th/0501033




[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)



Nucleon EM Form Factors: A Précis



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)



Nucleon EM Form Factors: A Précis



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)



Nucleon EM Form Factors: A Précis

Cloët, *et al.*:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

(Oettel, Hellstern, Alkofer, Reinhardt: [nucl-th/9805054](#))



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- But is that good?



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- But is that good?
 - Cloudy Bag: $\delta M_+^{\pi\text{-loop}} = -300$ to -400 MeV!



Nucleon EM Form Factors: A Précis

Cloët, *et al.*:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- But is that good?
 - Cloudy Bag: $\delta M_+^{\pi\text{-loop}} = -300$ to -400 MeV!
- Critical to anticipate pion cloud effects


Roberts, Tandy, Thomas, *et al.*, nu-th/02010084




Faddeev equation

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Faddeev equation




Faddeev equation



- Linear, Homogeneous Matrix equation
 - Yields *wave function* (Poincaré Covariant Faddeev Amplitude) that describes quark-diquark relative motion within the nucleon
- Scalar and Axial-Vector Diquarks ... In Nucleon's Rest Frame Amplitude has ... *s*–, *p*– & *d*–wave correlations



Diquark correlations



Diquark correlations

- Same interaction that describes mesons also generates three coloured quark-quark correlations: blue-red, blue-green, green-red


- Confined ... Does not escape from within baryon.

- Scalar is isosinglet, Axial-vector is isotriplet

- DSE and lattice-QCD

$$m_{[ud]_{0+}} = 0.74 - 0.82$$

$$m_{(uu)_{1+}} = m_{(ud)_{1+}} = m_{(dd)_{1+}} = 0.95 - 1.02$$



Nucleon-Photon Vertex

[First](#)[Contents](#)[Back](#)[Conclusion](#)

M. Oettel, M. Pichowsky
and L. von Smekal, nu-th/9909082

6 terms . . .

Nucleon-Photon Vertex

constructed systematically . . . current conserved automatically
for on-shell nucleons described by Faddeev Amplitude



[First](#)

[Contents](#)







[Back](#)

[Conclusion](#)


6 terms ...

Nucleon-Photon Vertex

constructed systematically ... current conserved automatically
for on-shell nucleons described by Faddeev Amplitude




Form Factor Ratio: GE/GM



Form Factor Ratio:


GE/GM

● Combine these elements ...



● Combine these elements ...


● Dressed-Quark Core



● Combine these elements ...


● Dressed-Quark Core

● *Ward-Takahashi*
Identity preserving
current




● Combine these elements ...

- Dressed-Quark Core
- *Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution



● Combine these elements ...

- Dressed-Quark Core
- *Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution




● Combine these elements ...

● Dressed-Quark Core

● *Ward-Takahashi*
Identity preserving
current


● Anticipate and
Estimate Pion
Cloud's Contribution

● All parameters fixed in
other applications ... **Not** varied.



● Combine these elements ...

- Dressed-Quark Core
- *Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution




● All parameters fixed in
other applications ... **Not** varied.

- Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$



- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution




- All parameters fixed in
other applications ... **Not** varied.

- Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$
- Correlations in Faddeev amplitude – quark orbital
angular momentum – essential to that agreement



- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution



- All parameters fixed in
other applications ... **Not** varied.

- Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$
- Correlations in Faddeev amplitude – quark orbital
angular momentum – essential to that agreement
- Predict Zero at $Q^2 \approx 6.5 \text{ GeV}^2$




Hall-A Neutron FF

- Preliminary result
E02-013 - Wojtsekhowski & Cates & Liyanage



- Preliminary result
- E02-013 - Wojtsekhowski & Cates & Liyanage



Improved current

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Improved current

- Composite axial-vector diquark correlation
 - Electromagnetic current can be complicated
 - Limited constraints on large- Q^2 behaviour



Improved current

- Composite axial-vector diquark correlation
 - Electromagnetic current can be complicated
 - Limited constraints on large- Q^2 behaviour
 - Improved performance of code
 - Implemented corrections so that large- Q^2 behaviour of form factors could be reliably calculated
 - Exposed two weaknesses in rudimentary *Ansatz*



Improved current

- Composite axial-vector diquark correlation
 - Improved performance of code
 - Implemented corrections so that large- Q^2 behaviour of form factors could be reliably calculated
 - Exposed two weaknesses in rudimentary *Ansatz*
 - Diquark effectively pointlike to hard probe
 - Didn't account for diquark being off-shell in recoil




Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%



Improved current


- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%


● Proton – zero shifted:
 $6.5 \rightarrow 8.0 \text{ GeV}^2$



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%

- **Proton** – zero shifted:
 $6.5 \rightarrow 8.0 \text{ GeV}^2$



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%


● **Proton** – zero shifted:

$$6.5 \rightarrow 8.0 \text{ GeV}^2$$

Neutron – peak shifted:

$$7.5 \rightarrow 5.0 \text{ GeV}^2$$

& now predict zero
a little above 11 GeV^2



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%


● **Proton** – zero shifted:

$$6.5 \rightarrow 8.0 \text{ GeV}^2$$

● **Neutron** – peak shifted:

$$7.5 \rightarrow 5.0 \text{ GeV}^2$$

& now predict zero
a little above 11 GeV^2






Pion Cloud

F_1 – neutron

- Comparison between Faddeev equation result and Kelly's parametrisation




Pion Cloud

F_1 – neutron

- Comparison

between Faddeev
equation result
and Kelly's
parametrisation


- Faddeev equation
set-up to describe
dressed-quark
core



Pion Cloud

F_1 – neutron

- Comparison between Faddeev equation result and Kelly's parametrisation
- Faddeev equation set-up to describe dressed-quark core
- Pseudoscalar meson cloud (and related effects) significant for $Q^2 \lesssim 3 - 4 M_N^2$





Epilogue



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)



Epilogue



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)



Epilogue

- DCSB exists in QCD.





Epilogue

- DCSB exists in QCD.
- It is manifest in dressed propagators and vertices





Epilogue

- DCSB exists in QCD.
 - It is manifest in dressed propagators and vertices
 - It predicts, amongst other things, that
 - light current-quarks become heavy constituent-quarks
 - pseudoscalar mesons are unnaturally light
 - pseudoscalar mesons couple unnaturally strongly to light-quarks
 - pseudoscalar mesons couple unnaturally strongly to the lightest baryons





Epilogue

- DCSB exists in QCD.
 - It is manifest in dressed propagators and vertices
 - It predicts, amongst other things, that
 - light current-quarks become heavy constituent-quarks
 - pseudoscalar mesons are unnaturally light
 - pseudoscalar mesons couple unnaturally strongly to light-quarks
 - pseudoscalar mesons couple unnaturally strongly to the lightest baryons
 - It impacts dramatically upon observables.





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors
 - Explaining the high Q^2 behavior of the proton form factor ratio in the space-like region





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors
 - Explaining the high Q^2 behavior of the proton form factor ratio in the space-like region
 - Detailing broadly the role of two-photon exchange contributions





nothing!

Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors
 - Explaining the high Q^2 behavior of the proton form factor ratio in the space-like region
 - Detailing broadly the role of two-photon exchange contributions
 - Explaining relationship between parton properties on the light-front and rest frame structure of hadrons



1. Universal Truths
2. QCD's Challenges
3. Why?
4. Some Questions
5. Status
6. Dichotomy of the Pion
7. Pion Form Factors
8. Dressed-Quark Propagator
9. Frontiers of Nuclear Science
10. Hadrons
11. Bethe-Salpeter Kernel
12. Persistent Challenge
13. Radial Excitations
14. Pion FF
15. Calculated Pion FF
16. $r_\pi f_\pi$
17. Two-photon Couplings
18. Two-photon Results
19. Nucleon Challenge
20. Nucleon EM Form Factors
21. Faddeev equation
22. Diquark correlations
23. Nucleon-Photon Vertex
24. Form Factor Ratio
25. Hall-A Neutron FF
26. Improved current
27. Pion Cloud
28. Dyson-Schwinger Equations
29. Schwinger Functions
30. Quenched-QCD
31. Pions and Form Factors
32. Masses: Nucleon and Δ
33. DIS Pion



Dyson-Schwinger Equations



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: **Generating Tool for Perturbation Theory**
..... **Materially Reduces** Model Dependence



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: **Generating Tool for Perturbation Theory**
..... **Materially Reduces** Model Dependence
- **NonPerturbative, Continuum approach to QCD**



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
..... Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
..... Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
..... Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?
- ⇒ Understanding InfraRed (long-range)
..... behaviour of $\alpha_s(Q^2)$



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
..... Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?
- Method yields Schwinger Functions \equiv Propagators



Dyson-Schwinger Equations

- Well suited to Relativistic Quantum Field Theory
- Simplest level: Generating Tool for Perturbation Theory
..... Materially Reduces Model Dependence
- NonPerturbative, Continuum approach to QCD
 - Hadrons as Composites of Quarks and Gluons
 - Qualitative and Quantitative Importance of:
 - Dynamical Chiral Symmetry Breaking
 - Generation of fermion mass from *nothing*
 - Quark & Gluon Confinement
 - Coloured objects not detected, not detectable?

Cross-Sections built from Schwinger Functions



Schwinger Functions

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Schwinger Functions

- Solutions are Schwinger Functions
(Euclidean **Green** Functions)



Schwinger Functions

- Solutions are Schwinger Functions (Euclidean **Green** Functions)
- Not all are Schwinger functions are experimentally observable



Schwinger Functions

- Solutions are Schwinger Functions (Euclidean **Green** Functions)
- Not all are Schwinger functions are experimentally observable but ...
 - **all are** same VEVs measured in numerical simulations of lattice-regularised QCD
 - opportunity for comparisons at pre-experimental level ... cross-fertilisation



Schwinger Functions

- Solutions are Schwinger Functions (Euclidean **Green** Functions)
- Not all are Schwinger functions are experimentally observable but ...
 - **all are** same VEVs measured in numerical simulations of lattice-regularised QCD
 - opportunity for comparisons at pre-experimental level ... cross-fertilisation
- Proving fruitful.




Persistent Challenge

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Persistent Challenge


● Infinitely Many Coupled Equations





Persistent Challenge

- Infinitely Many Coupled Equations




- Coupling between equations **necessitates** truncation





Persistent Challenge

- Infinitely Many Coupled Equations




- Coupling between equations **necessitates** truncation
 - Weak coupling expansion \Rightarrow Perturbation Theory





Persistent Challenge

- Infinitely Many Coupled Equations




- Coupling between equations **necessitates** truncation
 - Weak coupling expansion \Rightarrow Perturbation Theory
Not useful for the nonperturbative problems in which we're interested




Quenched-QCD

Dressed-Quark Propagator

$M(p)$




$Z(p)$




2002

$M(p)$



$Z(p)$




“*data*” Quenched Lattice Meas.

– Bowman, Heller, Leinweber, Williams: [he-lat/0209129](https://arxiv.org/abs/he-lat/0209129)


2002

Dressed-Quark Propagator

$M(p)$



$Z(p)$




“*data*,” Quenched Lattice Meas.

– Bowman, Heller, Leinweber, Williams: [he-lat/0209129](#)
current-quark masses: 30 MeV, 50 MeV, 100 MeV




2002

$M(p)$



$Z(p)$




- “*data*,” Quenched Lattice Meas.
 - Bowman, Heller, Leinweber, Williams: [he-lat/0209129](#)
 - current-quark masses: 30 MeV, 50 MeV, 100 MeV
- Curves: Quenched DSE Cal.
 - Bhagwat, Pichowsky, Roberts, Tandy [nu-th/0304003](#)




2002

$M(p)$



$Z(p)$



“*data*,” Quenched Lattice Meas.

– Bowman, Heller, Leinweber, Williams: [he-lat/0209129](#)
current-quark masses: 30 MeV, 50 MeV, 100 MeV

Curves: Quenched DSE Cal.

– Bhagwat, Pichowsky, Roberts, Tandy [nu-th/0304003](#)

Linear extrapolation of lattice data to chiral limit is **inaccurate**



Radial Excitations & Lattice-QCD

McNeile and Michael
he-la/0607032



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Radial Excitations & Lattice-QCD

McNeile and Michael
he-la/0607032

- *When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.*



Radial Excitations & Lattice-QCD

McNeile and Michael
he-la/0607032

- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.
- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$
 $\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$
Diehl & Hiller
he-ph/0105194



Radial Excitations & Lattice-QCD

McNeile and Michael
he-lq/0607032

- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.

- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$


$$\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$$

Diehl & Hiller

he-ph/0105194

- Lattice-QCD check:
 $16^3 \times 32$,
 $a \sim 0.1 \text{ fm}$,
two-flavour, unquenched

$$\Rightarrow \frac{f_{\pi_1}}{f_\pi} = 0.078 (93)$$



Radial Excitations & Lattice-QCD

McNeile and Michael
he-lq/0607032

- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.

- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$

$$\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$$

Diehl & Hiller

he-ph/0105194


- Lattice-QCD check:

$$16^3 \times 32,$$

$$a \sim 0.1 \text{ fm},$$

two-flavour, unquenched

$$\Rightarrow \frac{f_{\pi_1}}{f_\pi} = 0.078 (93)$$



- Full ALPHA formulation is required to see suppression, because PCAC relation is at the heart of the conditions imposed for improvement (determining coefficients of irrelevant operators)



Radial Excitations & Lattice-QCD

McNeile and Michael
he-lq/0607032

- When we first heard about [this result] our first reaction was a combination of “that is remarkable” and “unbelievable”.

- CLEO: $\tau \rightarrow \pi(1300) + \nu_\tau$

$$\Rightarrow f_{\pi_1} < 8.4 \text{ MeV}$$

Diehl & Hiller

he-ph/0105194


- Lattice-QCD check:

$$16^3 \times 32,$$

$$a \sim 0.1 \text{ fm},$$

two-flavour, unquenched

$$\Rightarrow \frac{f_{\pi_1}}{f_\pi} = 0.078 (93)$$




- The suppression of f_{π_1} is a useful benchmark that can be used to tune and validate lattice QCD techniques that try to determine the properties of excited states mesons.




Calculated Transition Form Factor:

RGI Rainbow-Ladder

Höll, Krassnigg, Maris, *et al.*,
“Electromagnetic properties of ground and
excited state pseudoscalar mesons,”
nu-th/0503043



 $m_u(1 \text{ GeV})$
 $= m_d(1 \text{ GeV}) = 5.5 \text{ MeV}$




Calculated Transition Form Factor:

Höll, Krassnigg, Maris, *et al.*,

“Electromagnetic properties of ground and excited state pseudoscalar mesons,”

nu-th/0503043

RGI Rainbow-Ladder



- $m_u(1 \text{ GeV}) = m_d(1 \text{ GeV}) = 5.5 \text{ MeV}$
- Predicted UV-behaviour is abundantly clear
 - precise for $Q^2 > 120 \text{ GeV}^2$



Harry Lee

Pions and Form Factors



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Pions and Form Factors


- Dynamical coupled-channels model . . . Analyzed extensive JLab data . . . Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C **54**, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C **63**, 055201/1-13 (2001)



Pions and Form Factors

- Dynamical coupled-channels model ... Analyzed extensive JLab data ... Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C **54**, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C **63**, 055201/1-13 (2001)
- Pion cloud effects are large in the low Q^2 region.

Ratio of the M1 form factor in $\gamma N \rightarrow \Delta$ transition and proton dipole form factor G_D . Solid curve is $G_M^(Q^2)/G_D(Q^2)$ including pions; Dotted curve is $G_M(Q^2)/G_D(Q^2)$ without pions.*




Pions and Form Factors

- Dynamical coupled-channels model ... Analyzed extensive JLab data ... Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C **54**, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C **63**, 055201/1-13 (2001)
- Pion cloud effects are large in the low Q^2 region.

Ratio of the M1 form factor in $\gamma N \rightarrow \Delta$ transition and proton dipole form factor G_D . Solid curve is $G_M^*(Q^2)/G_D(Q^2)$ including pions; Dotted curve is $G_M(Q^2)/G_D(Q^2)$ without pions.

Quark Core

- Responsible for only 2/3 of result at small Q^2
- Dominant for $Q^2 > 2 - 3 \text{ GeV}^2$



Results: Nucleon and Δ Masses



Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV)
for the scalar and axial-vector
diquark correlations, fixed by
fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for
Set B – fitted mass was offset to allow for “ π -cloud” contributions

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.80	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

● $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$



Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV) for the scalar and axial-vector diquark correlations, fixed by fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for
Set B – fitted mass was offset to allow for “ π -cloud” contributions

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.80	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

• $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$

• Axial-vector diquark provides significant attraction



Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV) for the scalar and axial-vector diquark correlations, fixed by fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for **Set B** – fitted mass was offset to allow for “ π -cloud” contributions

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.80	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

• $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$

• **Constructive Interference**: 1^{++} -diquark + $\partial_\mu \pi$



Deep-inelastic scattering




Deep-inelastic scattering



- Looking for Quarks




Deep-inelastic scattering



● Looking for Quarks



Deep-inelastic scattering




● Looking for Quarks

● **Signature Experiment** for QCD:

Discovery of Quarks at SLAC



Deep-inelastic scattering



● Looking for Quarks

● **Signature Experiment** for QCD:

Discovery of Quarks at SLAC

● Cross-section: Interpreted as Measurement of Momentum-Fraction Prob. Distribution: $q(x)$, $g(x)$



Pion's valence quark distn



Pion's valence quark distn

- π is Two-Body System: “Easiest” Bound State in QCD
- However, NO π Targets!



Pion's valence quark distn



- π is Two-Body System: “Easiest” Bound State in QCD
- However, NO π Targets!
- Existing Measurement Inferred from Drell-Yan:
$$\pi N \rightarrow \mu^+ \mu^- X$$




Pion's valence quark distn

- π is Two-Body System: “Easiest” Bound State in QCD
- However, NO π Targets!
- Existing Measurement Inferred from Drell-Yan:
 $\pi N \rightarrow \mu^+ \mu^- X$
- Proposal (Holt & Reimer, ANL, nu-ex/0010004)


$e^-_{5\text{GeV}} - p_{25\text{GeV}}$ Collider \rightarrow Accurate “Measurement”



Handbag diagrams



Handbag diagrams



$$W_{\mu\nu}(q; P) = \frac{1}{2\pi} \text{Im} [T_{\mu\nu}^+(q; P) + T_{\mu\nu}^-(q; P)]$$


$$T_{\mu\nu}^+(q, P) = \text{tr} \int \frac{d^4 k}{(2\pi)^4} \tau_- \bar{\Gamma}_\pi(k_{-\frac{1}{2}}; -P) S(k_{-0}) ieQ\Gamma_\nu(k_{-0}, k) \\ \times S(k) ieQ\Gamma_\mu(k, k_{-0}) S(k_{-0}) \tau_+ \Gamma_\pi(k_{-\frac{1}{2}}; P) S(k_{--})$$



Handbag diagrams

Bjorken Limit: $q^2 \rightarrow \infty$, $P \cdot q \rightarrow -\infty$
 but $x := -\frac{q^2}{2P \cdot q}$ fixed.

Numerous algebraic simplifications



$$W_{\mu\nu}(q; P) = \frac{1}{2\pi} \text{Im} [T_{\mu\nu}^+(q; P) + T_{\mu\nu}^-(q; P)]$$

$$T_{\mu\nu}^+(q, P) = \text{tr} \int \frac{d^4 k}{(2\pi)^4} \tau_- \bar{\Gamma}_\pi(k_{-\frac{1}{2}}; -P) S(k_{-0}) ieQ \Gamma_\nu(k_{-0}, k) \\ \times S(k) ieQ \Gamma_\mu(k, k_{-0}) S(k_{-0}) \tau_+ \Gamma_\pi(k_{-\frac{1}{2}}; P) S(k_{--})$$



Extant theory vs. experiment

K. Wijersooriya, P. Reimer and R. Holt,
nu-ex/0509012 ... Phys. Rev. C (Rapid)

